

Protocol of NFkB-BLA ME-180 Cell-based Assay for High-throughput Screening

DOCUMENT: NFkB-BLA_TOX21_SLP_Version1.0
TITLE: Protocol of NFkB-BLA ME-180 Cell-based Assay for High-throughput Screening

ASSAY REFERENCES:

Assay Target	Cell Lines	Species	Tissue of Origin	Assay Readout	Assay Provider	Toxicity Pathway
Nuclear Factor Kappa Beta (Endogenous)	ME-180	Human	Cervical carcinoma	Beta-lactamase reporter	Invitrogen	Stress response

QUALITY CONTROL PRECAUTIONS:

1. -The cells should not be grown more than 80-85% confluence
2. -The cell performance is affected is they are more confluent
3. -Do not leave cells in Trypsin for more than 5 min at RT
4. -handle 1536 well plate black clear bottom plates carefully by sides

MATERIALS and INSTRUMENTS:

Supplies/Medium/Reagent	Manufacturer	Vender/Catalog Number
-DMEM+Glutamax	-Invitrogen	-10569
-Dialyzed FBS	-Invitrogen	-26400
-NEAA	-Invitrogen	-11140
-HEPES	-Invitrogen	-15630
-Sodium Pyruvate	-Invitrogen	-11360
-Opti-MEM	-Invitrogen	-11058
-Penicillin Streptomycine	-Invitrogen	-15140
-Multidrop	-Thermofisher	-
-BiorapTR	-Beckman Coulter	-
-Envision	-Perkin Elmer	-
-LiveBLazer B/G FRET substrate	-Invitrogen	-K1030
-Blasticidin	-Invitrogen	-R21001

-0.25% Trypsin-EDTA	-Invitrogen	-25300
---------------------	-------------	--------

PROCEDURE:

1. Cell handling:

1.1. Media Required:

Component	Growth Medium	Assay Medium	Thaw Medium	Freezing Medium
-Recovery Cell Freezing Medium	-	-	-	-100%
-DMEM+ Glutamax	-90%	-	-90%	-
-OptiMEM	-	-99.5%	-	-
-Dialyzed FBS	-10%	-0.5%	-10%	-
-NEAA	-0.1 mM	-0.1 mM	-0.1 mM	-
-HEPES	-25 mM	-	-25 mM	-
-Sodium Pyruvate	-	-1 mM	-	-
-Penicillin- Streptomycin	-1%	-1%	-1%	-
-Blasticidin	-5 ug/ml	-	-	-
-	-	-	-	-

1.2. Thawing method

1.2.1 --Place 14 mL of pre-warmed thaw medium into a 15 ml conical tube

1.2.2 -Remove the vial of cells to be thawed from liquid nitrogen and thaw rapidly by placing at 37°C in a water bath with gentle agitation for 1-2 minutes. Do not submerge vial in water.

1.2.3 -Mix the entire content of the vial to 14 ml of pre-warmed medium and centrifuge to remove DMSO

1.2.4 -Discard the supernatant and transfer the precipitated cells to T175 flask using 30 ml thawing medium

1.3. Propagation method

1.3.1 -Detach the cells from the flask using 0.5 % Trypsin

1.3.2 -The cells are re-seeded in T-175 flask at 3-4 million

2. Assay Protocol

2.1 -Spin down the cells after rinsing the cells with DPBS and trypsinizing

2.2 -Resuspend the pellet with assay medium followed by filtering through cell strainer and adjust the required cell density

2.3 -Plate the cells in black-clear bottom 1536 well plate at 2000/well/6uL for agonist mode through 8 tip Multidrop plate dispenser

2.4 -Incubate for Overnight hrs at 37°C / 99% Humidity / 5% CO2

- 2.5 -Transfer 23nL of compounds from the library collection and positive control to the assay plates through pintool
 2.6 -Incubate for 5 hrs at 37°C / 99% Humidity / 5% CO2
 2.7 -Add 1uL of CCF4 (FRET Substrate) dye using a single tip plate dispenser (Bioraptr)
 2.8 --Incubate at room temperature for 2 hrs in dark
 2.9 -Read the fluorescence intensity through Envision plate reader using Beta-Lactamase protocol optimized for this cell type
 2.10 -Add 3 uL of Cell Titer Glo and Incubate at room temperature for 0.5 hrs in dark
 2.11 -Read on ViewLux protocol optimized for this cell type

3. Assay Performance

NFkB-bla Agonist (TNFa)	Online Validation (Mean ± SD)	Online Validation Viability (Mean ± SD)
EC50	0.107 ± 0.047 ng/mL (n = 27)	
S/B	7.88 ± 0.36	76.72 ± 89.24
CV (%)	4.78 ± 0.85 * (n = 18)	3.70 ± 0.23 * (n = 18)
Z'	0.86 ± 0.03	0.88 ± 0.04

* CV values shown represent average of DMSO plates and low concentration plates only.